



King's Research Portal

DOI:

[10.1002/eat.23158](https://doi.org/10.1002/eat.23158)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Huebel, C., Yilmaz, Z., Schaumberg, K., Breithaupt, L., Hunjan, A. K., García-gonzález, J., O'Reilly, P. F., Bulik, C. M., & Breen, G. D. (2019). Body composition in anorexia nervosa: meta-analysis and meta-regression of cross-sectional and longitudinal studies. *The International journal of eating disorders*, 52(11), 1205-1223. <https://doi.org/10.1002/eat.23158>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Body composition in anorexia nervosa: meta-analysis and meta-regression of cross-sectional and longitudinal studies

Christopher Hübel^{1,2,3}, Zeynep Yilmaz^{4,5}, Katherine E. Schaumberg^{4,6}, Lauren Breithaupt^{3,7,8}, Avina Hunjan^{1,2}, Judit García-González⁹, Paul F. O'Reilly¹, Cynthia M. Bulik^{3,4,10}, Gerome Breen^{1,2}

¹Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, UK

²UK National Institute for Health Research (NIHR) Biomedical Research Centre, South London and Maudsley Hospital, London, UK

³Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

⁴Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

⁵Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

⁶Department of Psychiatry, University of Wisconsin - Madison, Madison, WI, USA

⁷Eating Disorders Clinical and Research Program, Massachusetts General Hospital, Boston, USA

⁸Department of Psychiatry, Harvard Medical School, Boston, USA (LB)

⁹School of Biological and Chemical Sciences, Queen Mary, University of London, London, UK

¹⁰Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

PubMed indexing: Hübel, Yilmaz, Schaumberg, Breithaupt, Hunjan, García-González,
O'Reilly, Bulik, Breen

Correspondence to: Dr. Breen, Social, Genetic and Developmental Psychiatry Centre,
Institute of Psychiatry, Psychology and Neuroscience - PO80, De Crespigny Park, Denmark
Hill, London, United Kingdom, SE5 8AF; e-mail: gerome.breen@kcl.ac.uk; telephone:
+442078480409

Running head: Body composition in anorexia nervosa

Keywords: Weight restoration; body fat percentage; fat-free mass; lean mass; bone;
restricting; binge-eating/purging; long-term follow-up; BIA; bioelectrical impedance
analysis; DXA; dual-energy X-ray absorptiometry; insulin; estradiol; thyroid

Abbreviations: AN, anorexia nervosa; BIA, bioelectrical impedance analysis; DSM,
Diagnostic and Statistical Manual of Mental Disorders; DXA, dual-energy X-ray
absorptiometry; fT₃, free triiodothyronine; fT₄, free thyroxine; HDL, high-density lipoprotein;
IBW, ideal body weight; ICD, International Classification of Diseases, MRI, magnetic
resonance imaging; TSH, thyroid-stimulating hormone

Registry number: PROSPERO 2018 CRD42018105338

Abstract (236/250)

Objective: Clinically, anorexia nervosa (AN) presents with altered body composition. We quantified these alterations and evaluated their relationships with metabolites and hormones in AN patients longitudinally.

Method: In accordance with PRISMA guidelines, we conducted 88 meta-analyses on 60 samples published 1996-2018, comparing up to 2,294 pre-treatment, post-treatment, and weight-recovered AN patients with up to 1,848 controls. Primary outcomes were fat mass, fat-free mass, body fat percentage, and their regional distribution. Secondary outcomes were bone mineral density, metabolites, and hormones. Meta-regressions examined relationships among those measures and moderators.

Results: Pre-treatment AN patients evidenced 50% lower fat mass (MD: -9.05 kg, CI 95%: -10.02, -8.07, $Q = 9.33 \times 10^{-73}$) and 4.96 kg (CI 95%: -5.84, -4.07, $Q = 9.30 \times 10^{-27}$) lower fat-free mass, with fat mass preferentially stored in the trunk region during early weight restoration (4.2%, CI 95%: -2.1, -6.2, $Q = 2.30 \times 10^{-4}$). While the majority of traits returned to levels seen in healthy controls after weight restoration, fat-free mass (MD: -1.82 kg, CI 95%: -2.57, -1.08, $Q = 5.41 \times 10^{-6}$) and bone mineral density (MD: -0.10 kg, CI 95%: -0.18, -0.03, $Q = 0.01$) remained significantly altered.

Discussion: Body composition is markedly altered in AN, warranting research into these phenotypes as clinical risk or relapse predictors. Notably, the long term altered levels of fat-free mass and bone mineral density suggest that these parameters should be investigated as potential AN trait markers.

Introduction

Anorexia nervosa (AN) has one of the highest mortality rates of all psychiatric disorders (Chesney, Goodwin, & Fazel, 2014). Clinical observations show altered body composition (El Ghoch, Calugi, Lamborghini, & Dalle Grave, 2014; Solmi et al., 2016) accompanied by elevated cholesterol (Hussain et al., in press) and greater insulin sensitivity (Ilyas et al., 2018). However, conclusions are limited by small sample sizes and consequent mixed findings.

Molecular genetic studies have revealed that individuals with AN carry genetic variants that increase their liability to AN and concurrently predispose to lower body fat percentage, lower fasting insulin, and higher high-density lipoprotein (HDL) cholesterol concentrations, suggesting that metabolic factors may play an etiological role (Duncan et al., 2017; Watson et al., 2018). Additionally, longitudinal investigations of a British birth cohort showed that girls who develop AN later in life are already underweight at the age of 4 years when compared to healthy children (Yilmaz, Gottfredson, Zerwas, Bulik, & Micali, in press), adding evidence for a developmental component.

A systematic review showed that adolescents and adults differently lose fat tissue when affected by AN, with adolescents losing more central fat tissue and adults more peripheral fat tissue. During weight recovery, individuals with AN show emergent central adiposity which typically attenuates over time (El Ghoch, Calugi, et al., 2014). These clinical and genetic findings encourage the meta-analytic reassessment of the role of body composition traits, such as fat mass and fat-free mass, their regional distribution, and their changes associated with weight restoration and long-term weight recovery in AN.

The goals of these meta-analyses were to (1) replicate findings from the systematic review on fat mass; (2) extend the observations by quantifying them; (3) include fat-free mass and (4) bone mineral content and density; (5) investigate their associations with each other;

and (6) if possible, relate findings to secondary outcomes, such as metabolic and hormonal parameters. This analytical approach is aimed at understanding the potential associations between these factors that are known to be physiologically interrelated. A thorough and rigorous examination of body composition and related laboratory parameters in individuals with AN could elucidate some of the physiological changes associated with this serious disorder, which could lead to more effective medical management, monitoring, and treatment approaches.

Methods

Search strategy

Our meta-analysis was conducted according to PRISMA guidelines (Moher, Liberati, Tetzlaff, Altman, & PRISMA Group, 2009) and pre-registered (PROSPERO 2018 CRD42018105338) with no changes to the protocol. We conducted a literature search from 15.06.2018 until 01.07.2018 using the electronic database PubMed with a time limitation starting with articles published after 01.01.1994—marking the introduction of the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV; American Psychiatric Association, 2013)—until 31.06.2018. We used key search terms including "anorexia nervosa" AND ("body composition" OR "body fat" OR "fat mass" OR "body fat percentage" OR DXA OR BIA OR "fat free mass" OR "lean mass"). The search was repeated by co-authors to avoid selection bias. Furthermore, we screened the references of published articles and reviews. Our search results including the selection process are presented in Supporting Information Figure S1 according to PRISMA guidelines.

Selection criteria

Our inclusion criteria were as follows:

- a. Studies investigating humans only
- b. Any age group
- c. No sample overlap
- d. Observational cross-sectional or longitudinal studies or randomised-controlled trials
- e. Clinical diagnoses of AN according to the Diagnostic and Statistical Manual of Mental Disorders (DSM) IV, 5, or their revisions (American Psychiatric Association, 2013), or International Classification of Diseases version 10 (ICD-10) (World Health Organization, 1992)
- f. Investigation of body composition by dual-energy X-ray absorptiometry (DXA) (Bredella et al., 2010, 2013), bioelectrical impedance analysis (BIA) (Bonaccorsi et al., 2012; Mattar et al., 2011), dual photon absorptiometry (DPA), or magnetic resonance imaging (MRI) (Mayer et al., 2005).
- g. Published after 01.01.1994 (the year that DSM-IV was introduced)
- h. The study includes a control group or comparison group
- i. Publications in any language which could be translated by the research team: English, German, Swedish, Danish, Spanish

In case of multiple publications deriving from the same study population, we selected the articles reporting either the largest or the most recent data set. In case of conflict between these two criteria, large sample size was prioritized.

142 **Data extraction**

143 We extracted the following information from every identified study using a
144 standardised data extraction sheet:

- 145 a. Author, publication year
- 146 b. Country
- 147 c. Sample sizes including gender and age
- 148 d. Setting: inpatient or outpatient
- 149 e. Original longitudinal or cross-sectional design
- 150 f. Follow-up period if longitudinal
- 151 g. Diagnostic criteria: DSM-IV, DSM-IV-TR, DSM-5, or ICD-10
- 152 h. Participant screening and exclusion criteria
- 153 i. Number of cases: AN pre-treatment, post-treatment (ANpost), recovered from AN
154 (ANrec)
- 155 j. Subtype of AN: restricting (R), binge-eating/purging (BP)
- 156 k. Number of controls (CO)
- 157 l. Primary outcome variables of body composition: fat mass (FM), fat-free mass (FFM),
158 body fat percentage (BF%), and their regional distribution
- 159 m. Secondary outcome variables, which were reported by at least four studies additional
160 to primary outcomes: bone mineral density (BMD), glucose, insulin, ghrelin,
161 adiponectin, leptin, insulin-like growth factor (IGF-1), estradiol, testosterone, cortisol,
162 thyroid-stimulating hormone (TSH), free triiodothyronine (fT₃), free thyroxine (fT₄)
- 163 n. Covariates used in original analysis
- 164 o. Fasting and fasting duration
- 165 p. Blood sample: serum, plasma, or unspecified
- 166 q. Medication and contraceptives

- r. Psychological and additional treatments
- s. Outcome was a secondary or primary outcome in the original study
- t. Duration of illness
- u. Age at diagnosis/onset
- v. Age at menarche
- w. Percentage of AN cases with amenorrhea and duration of amenorrhea

Quality of study assessment (Newcastle-Ottawa Scale)

We used the Newcastle-Ottawa Scale (NOS) to assess the quality of studies (Wells et al., 2009). For the observational studies, low quality was defined as Newcastle–Ottawa Scale score ≤ 8.0 and high quality as score > 8.0 (maximum score 9).

Meta-analysis

Inverse variance-weighted meta-analyses were conducted using the statistical package ‘meta’ and ‘metafor’ in the open source software R v3.5.1 (r-project.org). We used additional formulas to calculate missing values (Hozo, Djulbegovic, & Hozo, 2005; Luo, Wan, Liu, & Tong, 2018; Wan, Wang, Liu, & Tong, 2014). Mean differences (MDs) between clinical and control groups were the effect sizes. A random-effects model, which assumes that the heterogeneity in the differences between clinical and control groups is due to both within-study and between-study variation, was used as we anticipated differences in procedures and study populations between studies. We used a restricted maximum-likelihood estimator (REML) to calculate the heterogeneity. For the analysis of subtypes, post-treatment, and weight-recovered AN patients, the control groups from the acutely-ill/pre-treatment analysis were reused because (1) control groups were not measured repeatedly; and (2) none of the studies had separate control groups for each subtype analysis. Although some studies

included covariates in their statistical analysis (Bratland-Sanda et al., 2010; Bredella et al., 2008; Dellava, Policastro, & Hoffman, 2009; DiVasta et al., 2007; Fernández-Soto, González-Jiménez, Chamorro-Fernández, & Leyva-Martínez, 2013; Haas et al., 2005; Karlsson, Weigall, Duan, & Seeman, 2000; Kosmiski, Schmiede, Mascolo, Gaudiani, & Mehler, 2014; Maimoun et al., 2018; Nakahara et al., 2007; Schneider et al., 1998), we only used raw values without including study-specific covariates to increase comparability across individual studies. Weight recovery was defined in accordance with DSM-IV and DSM-5 criteria with BMI >18.5 kg/m² or >90% ideal body weight (IBW). To correct our primary analysis for multiple testing, False Discovery Rate-adjusted *Q values* were calculated (Benjamini & Hochberg, 1995).

Publication bias or small study effects

Meta-analyses should diminish both bias and uncertainty of the overall estimate; however, this is not always the case as published studies may represent a biased selection of evidence (Nieminen, Rucker, Miettunen, Carpenter, & Schumacher, 2007). To detect publication bias, we inspected funnel plots visually (Egger, Smith, Schneider, & Minder, 1997) and quantified their asymmetry using a weighted linear regression by Thompson and Sharp—which allows for between-study heterogeneity (Thompson & Sharp, 1999). The significance of this test ($p < 0.05$) indicates asymmetric funnel plots and suggests publication bias or small study effects. In this case, we fitted a Copas selection model to adjust our meta-analysis for these effects.

Copas selection model

The Copas selection model to correct for potential selection bias comprises: (1) a model for the meta-analyzed effect; and (2) a model estimating the probability that study k is selected for publication. The correlation parameter ρ between these two components indicates the extent of publication bias with larger correlations suggesting more extreme effects have been selected for publication (Copas, 1999; Copas & Shi, 2000, 2001). We used the R package ‘metasens’ which fits the Copas selection model repeatedly over a grid of tuning parameters γ_0 and γ_1 and creates a standardised output: a) a funnel plot, b) a contour plot, c) a treatment effect plot, and d) a p value plot.

Meta-regression and stratified subgroup analysis

We detected extensive heterogeneity across the studies (Table 1). The heterogeneity observed in a meta-analysis can be partially systematic and related to clinical study-level variables (i.e., moderators), such as age or AN subtype. In secondary analyses, potential moderators were therefore explored in stratified subgroup analyses, such as AN subtype, or via meta-regression analyses, including mean age, the time period of follow-up for longitudinal studies, age at diagnosis, age at menarche, age at amenorrhea, duration of illness, percentage of amenorrhea in AN patients, percentage of medicated AN patients, percentage of individuals taking contraceptives, body composition measurement method, blood sample type, body composition parameters and their differences between cases and controls. Mixed effects models allow for within-study and between-study variation and were therefore the most appropriate model indicated by the large heterogeneity estimates detected in our primary meta-analyses. We used a restricted maximum-likelihood (REML) estimator to calculate the heterogeneity included in the R package ‘meta’.

Results

Results of the search and selection of studies

A total of 510 papers published between 1996 and 2018 were identified by our search terms, and 450 of them were excluded. No paper published 1994-1996 fulfilled the inclusion criteria and the most common reasons for exclusion were: (1) no control group ($n = 137$); (2) sample overlap ($n = 65$); and (3) no main outcome reported (i.e., body composition; $n = 45$). Detailed exclusion process is presented in Supporting Information Figure S1. Sixty published articles were included in our analysis (Agüera et al., 2015; Bachmann et al., 2014; Benninghoven, Raykowski, Solzbacher, Kunzendorf, & Jantschek, 2007; Bratland-Sanda et al., 2010; Bredella et al., 2012, 2008; Chudecka & Lubkowska, 2016; de Alvaro et al., 2007; Dellava et al., 2009; Delporte, Brichard, Hermans, Beguin, & Lambert, 2003; de Mateo Silleras et al., 2013; Diamanti et al., 2007; DiVasta et al., 2007, 2011; Dostálová, Sedláčková, Papezová, Nedvídková, & Haluzík, 2009; El Ghoch et al., 2012, 2015; El Ghoch, Calugi, Milanese, Bazzani, & Dalle Grave, 2017; El Ghoch, Milanese, et al., 2014; El Ghoch, Pourhassan, et al., 2017; Estour et al., 2017; Faje et al., 2014; Fazeli et al., 2010; Fernández-Soto et al., 2013; Galusca et al., 2015; Germain et al., 2010, 2007, 2016; Gniuli, Liverani, Capristo, Greco, & Mingrone, 2001; Grinspoon et al., 1996, 2001; Guo, Jiang, Liao, Liu, & He, 2013; Haas et al., 2018, 2005; Iacopino et al., 2003; Karczewska-Kupczewska et al., 2010; Karlsson et al., 2000; Kaváľková et al., 2012; Kerruish et al., 2002; Kirchengast & Huber, 2004; Konstantynowicz et al., 2011; Kosmiski et al., 2014; Maïmoun et al., 2018; Mayer et al., 2009, 2005; Mika, Herpertz-Dahlmann, Heer, & Holtkamp, 2004; Misra et al., 2013; Moreno, Djeddi, & Jaffrin, 2008; Mörkl et al., 2017; Nakahara et al., 2007; Nakai, Hamagaki, Takagi, Taniguchi, & Kurimoto, 1999; Prioletta et al., 2011; Rigaud, Boulier, Tallonneau, Brindisi, & Rozen, 2010; Scalfi, Marra, Caldara, Silvestri, & Contaldo, 1999; Scalfi et al., 2002; Schneider et al., 1998; Singhal et al., 2018; Tagami et al., 2004; Tanaka et

al., 2003; Weinbrenner et al., 2004), and we became aware of no additional unpublished
 samples after contacting study authors for additional or missing data (Supporting Information
 Table S1). The majority of studies focused on female cases and controls that were sampled
 consecutively in only 21 of 60 studies (Supporting Information Table S2) and aged between
 14.2 and 31.0 years (Supporting Information Figure S2). As such, two studies investigating
 male AN cases were excluded from the qualitative synthesis but are discussed briefly (El
 Ghoch, Calugi, et al., 2017; Misra et al., 2013). Three studies originated from Australasia, 37
 from Europe, 15 from North America, and five from Asia. Only 13 studies used the same
 method of ascertainment for cases and controls (Supporting Information Table S2). Twenty-
 eight studies investigated inpatients, eight outpatients, two a mixture of both, and 22 studies
 did not specify the recruitment or patient setting. All studies comprised collection of blood
 samples after a fasting period, whereas only six studies specified the fasting period (Bredella
 et al., 2012; DiVasta et al., 2011; Dostálová et al., 2009; Estour et al., 2017; Kaválková et al.,
 2012; Prioletta et al., 2011). One study did not specify whether analyses were performed
 using plasma or serum blood (Weinbrenner et al., 2004). Seventeen studies sampled regular
 menstruating participants during the follicular phase of their cycle (de Alvaro et al., 2007;
 Dostálová et al., 2009; Estour et al., 2017; Galusca et al., 2015; Germain et al., 2010, 2007,
 2016; Kaválková et al., 2012; Kirchengast & Huber, 2004; Mayer et al., 2005; Nakai et al.,
 1999; Prioletta et al., 2011; Weinbrenner et al., 2004), whereas 13 studies did not provide
 details about cycle phase (Bachmann et al., 2014; Bredella et al., 2012; Delporte et al., 2003;
 DiVasta et al., 2011; Fazeli et al., 2010; Fernández-Soto et al., 2013; Germain et al., 2010;
 Gniuli et al., 2001; Grinspoon et al., 1996; Guo et al., 2013; Haas et al., 2005; Karczewska-
 Kupczewska et al., 2010; Maïmoun et al., 2018; Mörk et al., 2017; Nakahara et al., 2007;
 Rigaud et al., 2010; Tagami et al., 2004; Tanaka et al., 2003). However, studies were retained
 to achieve the largest possible sample size, and—depending on data availability—meta-

regressions were fitted to investigate study characteristics as possible moderators. Originally, 39 studies were cross-sectional and 21 longitudinal (Supporting Information Table S1). However, three of the longitudinal studies were analysed cross-sectional in our meta-analysis due to missing data. No control group was repeatedly measured in any of the longitudinal studies.

Characteristics of the included studies

We performed four meta-analyses: 1) comparing 2,697 pre-treatment/acute-ill AN patients with 2,251 healthy controls; 2) comparing 722 post-treatment AN patients with 809 controls; 3) estimating the change in AN patients ($n = 722$) from pre- to post-treatment; and 4) comparing 398 weight-recovered individuals with AN with 660 healthy controls including samples with long-term follow-up. The pre-treatment AN group comprised 229 individuals suffering from the binge-eating/purging and 681 from the restricting subtype. The shortest follow-up period was 5.14 weeks, and the longest was 2 years (Supporting Information Table S1). Nineteen studies used BIA to assess body composition, 38 used DXA, and only three utilized MRI—considered to be the gold standard. Thirty of the 60 studies investigated body composition as a primary outcome, whereas it was a secondary outcome in the remaining studies. The percentage of AN patients with amenorrhea ranged from 43 to 100%, with ten studies not providing information on menstrual status (Agüera et al., 2015; Bachmann et al., 2014; Bredella et al., 2012; de Mateo Silleras et al., 2013; El Ghoch et al., 2012; Gniuli et al., 2001; Iacopino et al., 2003; Kirchengast & Huber, 2004; Schneider et al., 1998; Tagami et al., 2004; Tanaka et al., 2003). Thirty-five of 60 studies did not provide information on the medication status of AN patients, and 30 did not indicate whether oral contraceptives were used. In AN cases, the duration of illness was on average 52.59 months ($SD = 29.08$), the duration of amenorrhea 22.99 months ($SD = 18.27$), and the age at diagnosis 17.45 years (SD

= 3.01). Cases and controls were well matched for age (Supporting Information Figure S2) and, notably, we did not observe a difference in age at menarche (Supporting Information Figure S3) or height (Supporting Information Figure S6) between AN cases and controls.

Data and analyses results of meta-analyses and meta-regressions

Our results from the 88 meta-analyses show that a wide range of alterations in several key body composition and biochemical measures exist in AN cases compared with healthy controls (Figure 1/Supporting Information Figure S4). For 95% confidence intervals and Q values, heterogeneity estimates, and adjusted estimates due to estimated publication bias, see Table 1. Detailed forest plots showing each of the 88 meta-analyses are presented as Supporting Information Figures S5-92. No differences between restricting and binge-eating/purging subtype of AN were detected in our meta-analysis prior to treatment except for total body water (Supporting Information Table S3). Between-study heterogeneity (I^2) was observed in 62 meta-analyses and ranged from 52 to 99%, confirming our choice of a random-effects model. To investigate moderators implicated in heterogeneity, we performed 398 meta-regressions (Supporting Information Table S4-7). Eight meta-analyses showed funnel plot asymmetry, indicating small study effects. Therefore, we fitted Copas models to adjust for those effects and estimate the probable number of unpublished studies (Table 1 and Supporting Information Figure S93-100).

Primary outcomes: body composition

Anthropometrics

On average, pre-treatment AN cases had a 15.61 kg (CI 95%: -16.98, -14.24, $Q = 7.26 \times 10^{-109}$) lower body weight and were 0.01 m (CI 95%: -0.01, 0.00, $Q = 0.03$) shorter than healthy controls given their age ($\beta_{metareg} = -0.002$, $p = 0.04$; Supporting Information Table S4). After treatment, AN patients still weighed 4.92 kg (CI 95%: -8.03, -1.81, $Q = 1.92 \times 10^{-3}$) less than healthy controls.

Correspondingly, the pre-treatment BMI difference between AN cases and controls was -5.88 kg/m^2 (CI 95%: -6.31, -5.46, $Q = 9.06 \times 10^{-158}$), which reduced to -2.10 kg/m^2 (CI 95%: -2.53, -1.67, $p_{adjCopas} < 1.00 \times 10^{-4}$) after treatment as most patients gained on average 9.93 kg (CI 95%: 8.17, 11.68, $Q = 2.11 \times 10^{-27}$) during treatment. Post-treatment BMI was primarily accounted for by gains in fat mass ($\beta_{metareg} = 0.81$, $p = 7.03 \times 10^{-7}$) but not through fat-free mass (Supporting Information Table S5). After weight recovery, no statistically significant mean difference in BMI between AN cases and controls was detected.

Fat mass

The pre-treatment body composition of individuals with AN was significantly altered. Compared with healthy controls, AN cases had 9.05 kg (CI 95%: -10.02, -8.07, $Q = 9.33 \times 10^{-73}$) lower fat mass, corresponding to a 13.9% (CI 95%: -15.2, -12.6, $Q = 9.59 \times 10^{-99}$) lower complete body mass. This suggests that body fat was on average 50% lower than in healthy controls. Body fat percentage ($\beta_{metareg} = -134.53$, $p = 0.01$) and absolute fat mass ($\beta_{metareg} = -49.26$, $p = 1.65 \times 10^{-4}$) were associated with whole body bone mineral density of AN patients. Absolute fat mass was also associated with mean age at diagnosis ($\beta_{metareg} = -1.21$, $p = 2.42 \times 10^{-4}$; Supporting Information Table S4).

After treatment, AN patients had a 2.37 kg (CI 95%: -3.75, -0.98, $Q = 0.002$) lower fat mass which corresponded to 2.5% (CI 95%: -4.3, -0.7, $p_{adjCOPAS} = 0.006$) less total body mass compared with healthy controls. AN patients gained 6.39 kg (CI 95%: 5.13, 7.65, $Q = 3.07 \times 10^{-22}$) fat mass following treatment which corresponded to 10.4% (CI 95%: 7.96, 12.87, $Q = 6.25 \times 10^{-16}$) of total body mass. Post-treatment fat mass ($\beta_{metareg} = -0.23$, $p = 0.01$; Supporting Information Table S5) and gain in fat mass during treatment ($\beta_{metareg} = -0.20$, $p = 0.02$; Supporting Information Table S6) were negatively associated with the presence of amenorrhea. Following weight recovery, these values fully returned to levels seen in healthy controls.

Specifically, compared with healthy controls, AN patients had 3.51 kg (CI 95%: -4.58, -2.43, $Q = 8.07 \times 10^{-10}$) less trunk fat mass prior to treatment. In relative terms, however, AN patients had lower extremity body fat with 5.4% (CI 95%: -8.38, -2.43, $Q = 8.23 \times 10^{-4}$) less total body mass. The presence of amenorrhea was significantly associated with lower extremity fat mass ($\beta_{metareg} = 0.31$, $p = 0.04$; Supporting Information Table S4).

After treatment, AN patients showed a higher trunk body fat percentage than controls at 12.0% (CI 95%: 9.5, 14.4, $p_{adjCOPAS} < 1.00 \times 10^{-4}$) of total body mass. However, this finding was strongly influenced by publication bias with an estimated 52 unpublished studies. These results on body composition were not influenced by height as cases and controls showed no meaningful difference (i.e., 1 cm pre-treatment) or by age as meta-regressions were non-significant (Supporting Information Tables S4-7).

Fat-free mass

Overall, the fat-free mass in AN patients was 4.96 kg (CI 95%: -5.84, -4.07, $Q = 9.30 \times 10^{-27}$) lower pre-treatment than in controls, corresponding to 12.3% (CI 95%: 8.1, 16.5, $Q = 3.07 \times 10^{-8}$) higher proportion of total body mass. During treatment, AN patients gained 2.98 kg (CI 95%: 1.74, 4.22, $Q = 6.89 \times 10^{-6}$) fat-free mass, resulting in 1.82 kg (CI 95%: -2.57, -1.08, $Q = 5.41 \times 10^{-6}$) lower fat-free mass compared to controls. Yet, weight-recovered individuals with AN still showed 1.27 kg (CI 95%: -1.79, -0.75, $Q = 5.49 \times 10^{-6}$) lower fat-free mass than controls.

More specifically, pre-treatment fat-free mass of the extremities was 1.5% (CI 95%: -2.0, -1.0, $Q = 8.84 \times 10^{-9}$) less of total body mass. After treatment, no marked regional differences in fat-free mass were observed in AN patients. However, weight-recovered individuals with AN had 0.9% (CI 95%: -1.3, -0.5, $Q = 6.30 \times 10^{-5}$) lower trunk fat-free mass of total body mass than controls.

Before treatment, we observed a 393.95 kcal/day (CI 95%: -531.04, -256.86, $Q = 6.27 \times 10^{-8}$) lower resting energy expenditure and 4.65 L (CI 95%: -6.21, -3.10, $Q = 1.78 \times 10^{-8}$) less total body water in AN patients which persisted with 3.71 L (CI 95%: -7.07, -0.36, $Q = 0.05$) after treatment. Both were measured by BIA. However, resting energy expenditure was not corrected for fat-free mass or body mass in the original studies, limiting its interpretability. The association between total body water and fat-free mass could not be investigated as too few studies reported both outcomes concurrently.

Before treatment, only the amount of total body water was significantly different between individuals ($p_{\text{subgroup}} = 0.004$) suffering from the restricting (-5.21 L, CI 95%: -9.06, -1.36, $p_R = 8.04 \times 10^{-3}$, $k = 3$) or the binge-eating/purging subtype (-11.1 L, CI 95%: -12.04, -

10.16, $p_{BP} = 5.06 \times 10^{-119}$, $k = 1$; Supporting Information Table S3). However, this finding was limited by only one study investigating the binge-eating/purging subtype.

Secondary outcome: bone mineral measures

Bone mineral content and density

Compared with healthy controls, whole body bone mineral content in individuals with AN was 0.16 kg (CI 95%: -0.19, -0.12, $Q = 2.73 \times 10^{-20}$) lower pre-treatment and 0.09 kg (CI 95%: -0.13, -0.05, $Q = 1.63 \times 10^{-5}$) lower post-treatment. Weight-restored individuals with AN showed 0.10 kg (CI 95%: -0.18, -0.03, $Q = 0.01$) lower whole body bone mineral content compared to controls as they gained on average 0.05 kg (CI 95%: 0.02, 0.09, $Q = 0.01$) during treatment. The interpretability of these estimates is limited due to the insufficient follow-up time after weight recovery, exceeding six months in only two studies (Dellava et al., 2009; Karlsson et al., 2000). The pre-treatment whole body bone mineral content was associated with fat-free mass ($\beta_{metareg} = 0.02$, $p = 0.02$) and fat mass ($\beta_{metareg} = 0.05$, $p = 0.02$), as well as the difference in fat mass between AN patients and controls ($\beta_{metareg} = 0.04$, $p = 0.002$; Supporting Information Table S4). Accordingly, whole body pre-treatment bone mineral density was 0.03 g/cm² (CI 95%: -0.07, -0.00, $p_{adjCopas} = 0.02$) lower in AN, but our analysis showed a density comparable with healthy controls post-treatment. However, only two studies with 74 AN cases could be included in this analysis.

Before treatment, AN patients exhibited lower bone mineral density in several regions, including hip, lumbar spine, and femoral neck, with a few being likely to persist after weight recovery. These findings were associated with duration of illness, the age of AN cases, and differences in fat mass between cases and controls (Supporting Information: Secondary outcomes: detailed bone mineral measures and Table S4). Cases and controls in

our meta-analyses were age- and height-matched (Supporting Information Figure S2 & S6), therefore, these variables are unlikely to be associated with these results.

Secondary outcomes: metabolites and hormones

Exploratory results showed that fasting insulin and glucose concentrations were lower in AN patients compared with controls but not associated with fat or fat-free mass while lower leptin was associated with fat mass pre-treatment. After treatment, these measures returned to concentrations seen in healthy controls. Before treatment, thyroid hormones, cortisol, and IGF-1 were lower in AN patients and all three measures were associated with fat mass whereas higher cortisol in AN patients was associated with fat-free mass. For detailed results see Supporting Information: Secondary outcomes: metabolites and hormones.

Methodological moderators

We observed strong between-study heterogeneity (Table 1). To investigate how differences in study design, samples, and measurement methods may influence the primary and secondary outcomes, we performed an additional set of meta-regressions (Supporting Information Tables S4-7). The method of body composition measurement was associated with pre-treatment body fat percentage ($\beta_{DXA} = 3.34, p = 0.008$), fat-free mass ($\beta_{Isotope\ Dilution} = -6.18, p = 0.009$), and fat-free mass percentage ($\beta_{DXA} = -8.28, p = 0.01$) and post-treatment body fat percentage ($\beta_{DXA} = 6.39, p = 0.005$). Furthermore, femoral neck bone mineral density ($\beta_{Outpatient} = -0.12, p = 7.65 \times 10^{-4}$) significantly differed between inpatients and outpatients.

Discussion

Our primary meta-analyses showed marked alterations in body composition traits in patients with AN before and after treatment. Before treatment, all three major body compartments—fat, fat-free, and bone mass—showed significant reductions that were only partially restored after treatment. Our meta-analysis estimated ~50% lower body fat in AN patients which was mirrored by leptin concentrations (Perry & Shulman, 2018), both of which recovered with treatment. Significant differences were observed in body fat distribution post-treatment as body fat was primarily stored in the trunk. This distribution pattern may be due to increased insulin sensitivity observed in AN patients (Ilyas et al., 2018) potentially similar to observations in healthy individuals after short-term overfeeding (McLaughlin et al., 2016). We did not detect meaningful or statistically significant differences in body fat distribution in weight-restored patients, indicating potential redistribution occurring over longer term follow-up.

A new finding from our meta-analysis is that lower fat mass in AN was correlated with significantly low bone mineral content and density across the whole body. We speculate that the hormonal cross-talk between fat and bone tissue may be influencing this association (El Ghoch et al., 2016; Hawkes & Mostoufi-Moab, 2018), potentially mediated through greater bone marrow adipose tissue observed in AN (Fazeli & Klibanski, 2018; Suchacki & Cawthorn, 2018). Whole body bone mineral content remained low in weight-recovered individuals with AN. However, as only two studies followed patients for longer than six months (Dellava et al., 2009; Karlsson et al., 2000), the length of follow-up was insufficient to draw firm conclusions because bone mineral mass is slow to normalize. Future studies should be designed to capture long term changes. In men with AN, it has been reported that short-term weight restoration may normalize body composition patterns but could also lead to central adiposity (El Ghoch, Calugi, et al., 2017). However, sample sizes of reports of males

are very small. Additionally, in our analysis alterations in bone mineral mass did not affect the height of individuals with AN.

Another new finding in our meta-analysis is that we observed a 5 kg lower fat-free mass in AN patients which remained lower even after treatment and in weight-recovered AN patients, indicating that current treatment regimes may insufficiently target fat-free mass. Future studies should also assess muscle mass to identify the components of fat-free mass that are most associated with this reduction.

Our secondary outcomes—associations between detailed body composition and laboratory parameters in AN—were difficult to assess as only a few published studies reported both outcomes consistently. Most biochemical alterations were within the range of normal reference values. However, serious alterations can occur in certain individuals with AN that warrant vigilance by clinicians.

Overall, the meta-analyzed study sample was highly selected and biased as it comprised mostly European females aged between 14 and 31 years, emphasising the urgent need for studies including diverse ancestries, such as Asia, South and Central America, and Africa. Females and males differ in body composition and metabolic characteristics (Karastergiou & Fried, 2017; Link & Reue, 2017), underscoring the need for more studies on males with AN. Our study selection was limited by lack of control groups and underreported extensive sample overlap. Moreover, control groups were only measured at baseline in all longitudinal studies, failing to account for age- and growth-related variation, potentially inflating estimates. Furthermore, no clear-cut and consistent definition of recovery from AN was used across studies, contributing to heterogeneity (Murray, Loeb, & Le Grange, 2018). This underscores the necessity of developing standard definitions of remission and recovery in the eating disorders field (Bardone-Cone, Hunt, & Watson, 2018).

Methodologically, we observed effects of either BIA or DXA on the measurement of body composition in AN, questioning the comparability of the two methods. Larger, longitudinal validation studies comparing both methods with whole body MRI in eating disorders should be conducted. Additional factors influencing body composition and biochemical measures, such as menstrual cycle, diurnal changes, fasting, and pre-analytical procedures are summarised in Table 2 and should be carefully assessed in future studies (Hernandes, Barbas, & Dudzik, 2017).

Most importantly, blood is comprised of approximately 3,500 highly correlated and interacting proteins (hupo.org) (Schwenk et al., 2017) and 4,600 metabolites (serummetabolome.ca) (Psychogios et al., 2011), therefore, the measurement of single proteins, hormones or metabolites is ill-advised. Metabolomics, proteomics, and lipidomics can capture large amounts of information at adequate statistical power when used in large samples (Hernandes et al., 2017). Additionally, large epidemiological databases that have measured biomarkers in childhood, such as the Avon Longitudinal Study of Parents and Children (ALSPAC; Golding, Pembrey, Jones, & ALSPAC Study Team, 2001) and Generation R (Kooijman et al., 2016), should be harnessed to determine whether those who go on to develop AN show evidence for premorbid differences in body composition and biochemical parameters as has been observed for BMI by Yilmaz et al. (Yilmaz et al., in press).

Conclusions

Detailed measurement of body composition with simple methods, such as BIA or DXA, that offers additional information on bone tissue, may help refine our understanding of the nature of AN and its diagnosis. Our meta-analyses showed that all body compartments were markedly altered in AN. Individuals with AN presented with 50% lower fat mass and prolonged recovery periods for fat-free mass and bone mineral content. The core implication of body composition differences are alterations in metabolism, growth, and development. While results must be interpreted with caution given small samples, we found evidence indicating alterations in fasting insulin, thyroid, sex, and stress hormones in AN which appeared to partially normalize with weight gain and recovery. Large birth cohorts that collected information on eating disorders along with metabolomic information offer a rich and exciting opportunity for prospective investigations that add to our understanding of body composition and metabolic mechanisms in risk and maintenance of eating disorders.

Acknowledgments

Prof. Bulik acknowledges funding from the Swedish Research Council (VR Dnr: 538-2013-8864). Prof Bulik is supported by NIMH R21 MH115397. Dr. Yilmaz acknowledges grant support from NIH K01 MH109782. This study represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. No funder had a role in the design, analysis or writing of this article. PFO receives funding from the UK Medical Research Council (MR/N015746/1) and the Wellcome Trust (109863/Z/15/Z).

Conflict of interest

Dr. Bulik reports: Shire (Scientific Advisory Board member) and Pearson (author, royalty recipient) (unrelated to the content of this manuscript). Dr. Breen has received grant funding from and served as a consultant to Eli Lilly and has received honoraria from Illumina and has served on advisory boards for Otsuka (all unrelated to the content of this manuscript). The remaining authors declare that they have no conflict of interest.

Authors' contributions

CH, ZY, CMB and GB designed research; CH, ZY, KS, LB, AH, JGG conducted research; CH, ZY, KS, LB, AH, JGG provided essential materials; CH analyzed data or performed statistical analysis; CH, ZY, LB, KS, GB, CMB wrote paper; CH had primary responsibility for final content. All authors read and approved the final manuscript.

563 **Data availability**

564 All data and all scripts used for data analysis are available on

565 github.com/topherhuebel/metabcan

566

References

- Agüera, Z., Romero, X., Arcelus, J., Sánchez, I., Riesco, N., Jiménez-Murcia, S., ... Fernández-Aranda, F. (2015). changes in body composition in anorexia nervosa: Predictors of recovery and treatment outcome. *PloS One*, *10*(11), e0143012.
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). Washington, DC: American Psychiatric Association.
- Bachmann, K. N., Fazeli, P. K., Lawson, E. A., Russell, B. M., Riccio, A. D., Meenaghan, E., ... Miller, K. K. (2014). Comparison of hip geometry, strength, and estimated fracture risk in women with anorexia nervosa and overweight/obese women. *The Journal of Clinical Endocrinology and Metabolism*, *99*(12), 4664–4673.
- Bardone-Cone, A. M., Hunt, R. A., & Watson, H. J. (2018). An overview of conceptualizations of eating disorder recovery, recent findings, and future directions. *Current Psychiatry Reports*, *20*(9), 79.
- Benjamini, Y., & Hochberg, Y. (1995). controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B, Statistical Methodology*, *57*(1), 289–300.
- Benninghoven, D., Raykowski, L., Solzbacher, S., Kunzendorf, S., & Jantschek, G. (2007). Body images of patients with anorexia nervosa, bulimia nervosa and female control subjects: a comparison with male ideals of female attractiveness. *Body Image*, *4*(1), 51–59.
- Bonaccorsi, G., Bassetti, A., Chiari, S., Dirindelli, P., Lorini, C., Menicalli, C., ... Martinetti, M. G. (2012). Body composition in subjects with anorexia nervosa: bioelectrical impedance analysis and dual-energy X-ray absorptiometry. *Eating and Weight Disorders*, *17*(4), e298–e303.
- Bratland-Sanda, S., Sundgot-Borgen, J., Rosenvinge, J. H., Rø, Ø., Hoffart, A., & Martinsen, E. W. (2010). Physical fitness, bone mineral density and associations with physical activity in females with longstanding eating disorders and non-clinical controls. *Journal of Sports Medicine and Physical Fitness*, *50*(3), 303–310.
- Bredella, M. A., Fazeli, P. K., Freedman, L. M., Calder, G., Lee, H., Rosen, C. J., & Klibanski, A.

- 594 (2012). Young women with cold-activated brown adipose tissue have higher bone mineral
595 density and lower Pref-1 than women without brown adipose tissue: a study in women with
596 anorexia nervosa, women recovered from anorexia nervosa, and normal-weight women. *The*
597 *Journal of Clinical Endocrinology and Metabolism*, 97(4), E584–E590.
- 598 Bredella, M. A., Ghomi, R. H., Thomas, B. J., Torriani, M., Brick, D. J., Gerweck, A. V., ... Miller,
599 K. K. (2010). Comparison of DXA and CT in the assessment of body composition in
600 premenopausal women with obesity and anorexia nervosa. *Obesity*, 18(11), 2227–2233.
- 601 Bredella, M. A., Gill, C. M., Keating, L. K., Torriani, M., Anderson, E. J., Punyanitya, M., ... Miller,
602 K. K. (2013). Assessment of abdominal fat compartments using DXA in premenopausal women
603 from anorexia nervosa to morbid obesity. *Obesity*, 21(12), 2458–2464.
- 604 Bredella, M. A., Misra, M., Miller, K. K., Madisch, I., Sarwar, A., Cheung, A., ... Gupta, R. (2008).
605 Distal radius in adolescent girls with anorexia nervosa: Trabecular structure analysis with high-
606 resolution flat-panel volume CT. *Radiology*, 249(3), 938–946.
- 607 Chesney, E., Goodwin, G. M., & Fazel, S. (2014). Risks of all-cause and suicide mortality in mental
608 disorders: A meta-review. *World Psychiatry*, 13(2), 153–160.
- 609 Chudecka, M., & Lubkowska, A. (2016). Thermal imaging of body surface temperature distribution in
610 women with anorexia nervosa. *European Eating Disorders Review*, 24(1), 57–61.
- 611 Copas, J. (1999). What works?: selectivity models and meta-analysis. *Journal of the Royal Statistical*
612 *Society*, 162(1), 95–109.
- 613 Copas, J., & Shi, J. Q. (2000). Meta-analysis, funnel plots and sensitivity analysis. *Biostatistics*, 1(3),
614 247–262.
- 615 Copas, J., & Shi, J. Q. (2001). A sensitivity analysis for publication bias in systematic reviews.
616 *Statistical Methods in Medical Research*, 10(4), 251–265.
- 617 de Alvaro, M. T. G., Muñoz-Calvo, M. T., Barrios, V., Martínez, G., Martos-Moreno, G. A.,
618 Hawkins, F., & Argente, J. (2007). Regional fat distribution in adolescents with anorexia
619 nervosa: effect of duration of malnutrition and weight recovery. *European Journal of*
620 *Endocrinology*, 157(4), 473–479.
- 621 Dellava, J. E., Policastro, P., & Hoffman, D. J. (2009). Energy metabolism and body composition in

- 622 long-term recovery from anorexia nervosa. *International Journal of Eating Disorders*, 42(5),
623 415–421.
- 624 Delporte, M. L., Brichard, S. M., Hermans, M. P., Beguin, C., & Lambert, M. (2003).
625 Hyperadiponectinaemia in anorexia nervosa. *Clinical Endocrinology*, 58(1), 22–29.
- 626 de Mateo Silleras, B., Redondo del Río, P., Camina Martín, A., Soto Célix, M., Alonso Torre, S. R.,
627 & Miján de la Torre, A. (2013). [Effect of refeeding on the body composition of females with
628 restrictive anorexia nervosa; anthropometry versus bioelectrical impedance]. *Nutricion*
629 *hospitalaria*, 28(5), 1717–1724.
- 630 Diamanti, A., Bizzarri, C., Gambarara, M., Calce, A., Montecchi, F., Cappa, M., ... Castro, M.
631 (2007). Bone mineral density in adolescent girls with early onset of anorexia nervosa. *Clinical*
632 *Nutrition*, 26(3), 329–334.
- 633 DiVasta, A. D., Beck, T. J., Petit, M. A., Feldman, H. A., LeBoff, M. S., & Gordon, C. M. (2007).
634 Bone cross-sectional geometry in adolescents and young women with anorexia nervosa: A hip
635 structural analysis study. *Osteoporosis International*, 18(6), 797–804.
- 636 DiVasta, A. D., Feldman, H. A., Brown, J. N., Giancaterino, C., Holick, M. F., & Gordon, C. M.
637 (2011). Bioavailability of vitamin D in malnourished adolescents with anorexia nervosa. *Journal*
638 *of Clinical Endocrinology and Metabolism*, 96(8), 2575–2580.
- 639 Dostálová, I., Sedláčková, D., Papezová, H., Nedvídková, J., & Haluzík, M. (2009). Serum visfatin
640 levels in patients with anorexia nervosa and bulimia nervosa. *Physiological Research*, 58(6),
641 903–907.
- 642 Duncan, L., Yilmaz, Z., Gaspar, H., Walters, R., Goldstein, J., Anttila, V., ... Bulik, C. M. (2017).
643 Significant locus and metabolic genetic correlations revealed in genome-wide association study
644 of anorexia nervosa. *American Journal of Psychiatry*, 174(9), 850–858.
- 645 Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a
646 simple, graphical test. *BMJ*, 315(7109), 629–634.
- 647 El Ghoch, M., Alberti, M., Milanese, C., Battistini, N. C., Pellegrini, M., Capelli, C., ... Dalle Grave,
648 R. (2012). Comparison between dual-energy X-ray absorptiometry and skinfolds thickness in
649 assessing body fat in anorexia nervosa before and after weight restoration. *Clinical Nutrition*,

- 650 31(6), 911–916.
- 651 El Ghoch, M., Calugi, S., Lamburghini, S., & Dalle Grave, R. (2014). Anorexia nervosa and body fat
652 distribution: A systematic review. *Nutrients*, 6(9), 3895–3912.
- 653 El Ghoch, M., Calugi, S., Milanese, C., Bazzani, P. V., & Dalle Grave, R. (2017). Body composition
654 in men with anorexia nervosa: Longitudinal study. *International Journal of Eating Disorders*,
655 50(7), 856–860.
- 656 El Ghoch, M., Gatti, D., Calugi, S., Viapiana, O., Bazzani, P. V., & Dalle Grave, R. (2016). The
657 association between weight gain/restoration and bone mineral density in adolescents with
658 anorexia nervosa: A systematic review. *Nutrients*, 8(12). <https://doi.org/10.3390/nu8120769>
- 659 El Ghoch, M., Milanese, C., Calugi, S., Müller, M. J., Pourhassan, M., Ruocco, A., & Dalle Grave, R.
660 (2015). Regional fat distribution in adolescent and adult females with anorexia nervosa: A
661 longitudinal study. *Clinical Nutrition*, 34(6), 1224–1232.
- 662 El Ghoch, M., Milanese, C., Calugi, S., Pellegrini, M., Battistini, N. C., & Dalle Grave, R. (2014).
663 Body composition, eating disorder psychopathology, and psychological distress in anorexia
664 nervosa: a longitudinal study. *American Journal of Clinical Nutrition*, 99(4), 771–778.
- 665 El Ghoch, M., Pourhassan, M., Milanese, C., Müller, M. J., Calugi, S., Bazzani, P. V., & Dalle Grave,
666 R. (2017). Changes in lean and skeletal muscle body mass in adult females with anorexia nervosa
667 before and after weight restoration. *Clinical Nutrition*, 36(1), 170–178.
- 668 Estour, B., Marouani, N., Sigaud, T., Lang, F., Fakra, E., Ling, Y., ... Germain, N. (2017).
669 Differentiating constitutional thinness from anorexia nervosa in DSM 5 era.
670 *Psychoneuroendocrinology*, 84, 94–100.
- 671 Faje, A. T., Fazeli, P. K., Miller, K. K., Katzman, D. K., Ebrahimi, S., Lee, H., ... Klibanski, A.
672 (2014). Fracture risk and areal bone mineral density in adolescent females with anorexia nervosa.
673 *International Journal of Eating Disorders*, 47(5), 458–466.
- 674 Fazeli, P. K., Bredella, M. A., Misra, M., Meenaghan, E., Rosen, C. J., Clemmons, D. R., ...
675 Klibanski, A. (2010). Preadipocyte factor-1 is associated with marrow adiposity and bone
676 mineral density in women with anorexia nervosa. *Journal of Clinical Endocrinology and*
677 *Metabolism*, 95(1), 407–413.

- 678 Fazeli, P. K., & Klibanski, A. (2018). The paradox of marrow adipose tissue in anorexia nervosa.
679 *Bone*. <https://doi.org/10.1016/j.bone.2018.02.013>
- 680 Fernández-Soto, M. L., González-Jiménez, A., Chamorro-Fernández, M., & Leyva-Martínez, S.
681 (2013). Clinical and hormonal variables related to bone mass loss in anorexia nervosa patients.
682 *Vitamins and Hormones*, 92, 259–269.
- 683 Galusca, B., Prevost, G., Germain, N., Dubuc, I., Ling, Y., Anouar, Y., ... Chartrel, N. (2015).
684 Neuropeptide Y and alpha-MSH circadian levels in two populations with low body weight:
685 Anorexia nervosa and constitutional thinness. *PloS One*, 10(3), e0122040.
- 686 Germain, N., Galusca, B., Grouselle, D., Frere, D., Billard, S., Epelbaum, J., & Estour, B. (2010).
687 Ghrelin and obestatin circadian levels differentiate bingeing-purging from restrictive anorexia
688 nervosa. *Journal of Clinical Endocrinology and Metabolism*, 95(6), 3057–3062.
- 689 Germain, N., Galusca, B., Le Roux, C. W., Bossu, C., Ghatei, M. A., Lang, F., ... Estour, B. (2007).
690 Constitutional thinness and lean anorexia nervosa display opposite concentrations of peptide YY,
691 glucagon-like peptide 1, ghrelin, and leptin. *American Journal of Clinical Nutrition*, 85(4), 967–
692 971.
- 693 Germain, N., Viltart, O., Loyens, A., Bruchet, C., Nadin, K., Wolowczuk, I., ... Galusca, B. (2016).
694 Interleukin-7 plasma levels in human differentiate anorexia nervosa, constitutional thinness and
695 healthy obesity. *PloS One*, 11(9), e0161890.
- 696 Gniuli, D., Liverani, E., Capristo, E., Greco, A. V., & Mingrone, G. (2001). Blunted glucose
697 metabolism in anorexia nervosa. *Metabolism: Clinical and Experimental*, 50(8), 876–881.
- 698 Golding, J., Pembrey, M., Jones, R., & ALSPAC Study Team. (2001). ALSPAC--the Avon
699 Longitudinal Study of Parents and Children. I. Study methodology. *Paediatric and Perinatal*
700 *Epidemiology*, 15(1), 74–87.
- 701 Grinspoon, S., Gulick, T., Askari, H., Landt, M., Lee, K., Anderson, E., ... Klibanski, A. (1996).
702 Serum leptin levels in women with anorexia nervosa. *Journal of Clinical Endocrinology and*
703 *Metabolism*, 81(11), 3861–3863.
- 704 Grinspoon, S., Thomas, L., Miller, K., Pitts, S., Herzog, D., & Klibanski, A. (2001). Changes in
705 regional fat redistribution and the effects of estrogen during spontaneous weight gain in women

- 706 with anorexia nervosa. *American Journal of Clinical Nutrition*, 73(5), 865–869.
- 707 Guo, L. J., Jiang, T. J., Liao, L., Liu, H., & He, H. B. (2013). Relationship between serum omentin-1
708 level and bone mineral density in girls with anorexia nervosa. *Journal of Endocrinological*
709 *Investigation*, 36(3), 190–194.
- 710 Haas, V., Kent, D., Kohn, M. R., Madden, S., Clarke, S., Briody, J., ... Gaskin, K. (2018). Incomplete
711 total body protein recovery in adolescent patients with anorexia nervosa. *American Journal of*
712 *Clinical Nutrition*, 107(3), 303–312.
- 713 Haas, V., Onur, S., Paul, T., Nutzinger, D. O., Bösby-Westphal, A., Hauer, M., ... Müller, M. J.
714 (2005). Leptin and body weight regulation in patients with anorexia nervosa before and during
715 weight recovery. *American Journal of Clinical Nutrition*, 81(4), 889–896.
- 716 Hawkes, C. P., & Mostoufi-Moab, S. (2018). Fat-bone interaction within the bone marrow milieu:
717 Impact on hematopoiesis and systemic energy metabolism. *Bone*.
718 <https://doi.org/10.1016/j.bone.2018.03.012>
- 719 Hernandez, V. V., Barbas, C., & Dudzik, D. (2017). A review of blood sample handling and pre-
720 processing for metabolomics studies. *Electrophoresis*, 38(18), 2232–2241.
- 721 Hozo, S. P., Djulbegovic, B., & Hozo, I. (2005). Estimating the mean and variance from the median,
722 range, and the size of a sample. *BMC Medical Research Methodology*, 5, 13.
- 723 Hussain, A. A., Hübel, C., Hindborg, M., Lindkvist, E., Kastrup, A., Yilmaz, Z., ... Sjögren, M. (in
724 press). Increased lipid and lipoprotein concentrations in anorexia nervosa: A systematic review
725 and meta-analysis. *International Journal of Eating Disorders*.
- 726 Iacopino, L., Siani, V., Melchiorri, G., Orlandi, C., De Luna, A., Cervelli, V., & Andreoli, A. (2003).
727 Body composition differences in adolescent female athletes and anorexic patients. *Acta*
728 *Diabetologica*, 40 Suppl 1, S180–S182.
- 729 Ilyas, A., Hübel, C., Stahl, D., Stadler, M., Ismail, K., Breen, G., ... Kan, C. (2018). The metabolic
730 underpinning of eating disorders: A systematic review and meta-analysis of insulin sensitivity.
731 *Molecular and Cellular Endocrinology*. <https://doi.org/10.1016/j.mce.2018.10.005>
- 732 Karastergiou, K., & Fried, S. K. (2017). Cellular mechanisms driving sex differences in adipose tissue
733 biology and body shape in humans and mouse models. *Advances in Experimental Medicine and*

- 734 *Biology*, 1043, 29–51.
- 735 Karczeńska-Kupczewska, M., Straczkowski, M., Adamska, A., Nikolajuk, A., Otziomek, E., Gorska,
 736 M., & Kowalska, I. (2010). Insulin sensitivity, metabolic flexibility, and serum adiponectin
 737 concentration in women with anorexia nervosa. *Metabolism: Clinical and Experimental*, 59(4),
 738 473–477.
- 739 Karlsson, M. K., Weigall, S. J., Duan, Y., & Seeman, E. (2000). Bone size and volumetric density in
 740 women with anorexia nervosa receiving estrogen replacement therapy and in women recovered
 741 from anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism*, 85(9), 3177–3182.
- 742 Kaváľková, P., Dostálová, I., Haluzíková, D., Trachta, P., Hanušová, V., Lacinová, Z., ... Haluzík, M.
 743 (2012). Preadipocyte factor-1 concentrations in patients with anorexia nervosa: the influence of
 744 partial realimentation. *Physiological Research*, 61(2), 153–159.
- 745 Kerruish, K. P., O'Connor, J., Humphries, I. R. J., Kohn, M. R., Clarke, S. D., Briody, J. N., ... Baur,
 746 L. A. (2002). Body composition in adolescents with anorexia nervosa. *American Journal of*
 747 *Clinical Nutrition*, 75(1), 31–37.
- 748 Kirchengast, S., & Huber, J. (2004). Body composition characteristics and fat distribution patterns in
 749 young infertile women. *Fertility and Sterility*, 81(3), 539–544.
- 750 Konstantynowicz, J., Abramowicz, P., Jamiolkowski, J., Kadziela-Olech, H., Bialokoz-Kalinowska,
 751 I., Kierus-Jankowska, K., ... Kaczmarek, M. (2011). Thigh circumference as a useful predictor
 752 of body fat in adolescent girls with anorexia nervosa. *Annals of Nutrition & Metabolism*, 58(3),
 753 181–187.
- 754 Kooijman, M. N., Kruithof, C. J., van Duijn, C. M., Duijts, L., Franco, O. H., van IJzendoorn, M. H.,
 755 ... Jaddoe, V. W. V. (2016). The Generation R Study: Design and cohort update 2017. *European*
 756 *Journal of Epidemiology*, 31(12), 1243–1264.
- 757 Kosmiski, L., Schmiege, S. J., Mascolo, M., Gaudiani, J., & Mehler, P. S. (2014). Chronic starvation
 758 secondary to anorexia nervosa is associated with an adaptive suppression of resting energy
 759 expenditure. *Journal of Clinical Endocrinology and Metabolism*, 99(3), 908–914.
- 760 Link, J. C., & Reue, K. (2017). Genetic Basis for Sex Differences in Obesity and Lipid Metabolism.
 761 *Annual Review of Nutrition*, 37, 225–245.

- 762 Luo, D., Wan, X., Liu, J., & Tong, T. (2018). Optimally estimating the sample mean from the sample
763 size, median, mid-range, and/or mid-quartile range. *Statistical Methods in Medical Research*,
764 27(6), 1785–1805.
- 765 Maïmoun, L., Guillaume, S., Lefebvre, P., Bertet, H., Seneque, M., Philibert, P., ... Sultan, C. (2018).
766 Effects of the two types of anorexia nervosa (binge eating/purging and restrictive) on bone
767 metabolism in female patients. *Clinical Endocrinology*, 88(6), 863–872.
- 768 Mattar, L., Godart, N., Melchior, J. C., Falissard, B., Kolta, S., Ringuenet, D., ... Pichard, C. (2011).
769 Underweight patients with anorexia nervosa: Comparison of bioelectrical impedance analysis
770 using five equations to dual X-ray absorptiometry. *Clinical Nutrition*, 30(6), 746–752.
- 771 Mayer, L. E. S., Klein, D. A., Black, E., Attia, E., Shen, W., Mao, X., ... Walsh, B. T. (2009).
772 Adipose tissue distribution after weight restoration and weight maintenance in women with
773 anorexia nervosa. *American Journal of Clinical Nutrition*, 90(5), 1132–1137.
- 774 Mayer, L. E. S., Walsh, B. T., Pierson, R. N., Jr, Heymsfield, S. B., Gallagher, D., Wang, J., ...
775 Glasofer, D. (2005). Body fat redistribution after weight gain in women with anorexia nervosa.
776 *American Journal of Clinical Nutrition*, 81(6), 1286–1291.
- 777 McLaughlin, T., Craig, C., Liu, L.-F., Perelman, D., Allister, C., Spielman, D., & Cushman, S. W.
778 (2016). Adipose cell size and regional fat deposition as predictors of metabolic response to
779 overfeeding in insulin-resistant and insulin-sensitive humans. *Diabetes*, 65(5), 1245–1254.
- 780 Mika, C., Herpertz-Dahlmann, B., Heer, M., & Holtkamp, K. (2004). Improvement of nutritional
781 status as assessed by multifrequency BIA during 15 weeks of refeeding in adolescent girls with
782 anorexia nervosa. *Journal of Nutrition*, 134(11), 3026–3030.
- 783 Misra, M., Katzman, D. K., Clarke, H., Snelgrove, D., Brigham, K., Miller, K. K., & Klibanski, A.
784 (2013). Hip structural analysis in adolescent boys with anorexia nervosa and controls. *Journal of*
785 *Clinical Endocrinology and Metabolism*, 98(7), 2952–2958.
- 786 Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group. (2009). Preferred reporting
787 items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal*
788 *Medicine*, 151(4), 264–269, W64.
- 789 Moreno, M. V., Djeddi, D. D., & Jaffrin, M. Y. (2008). Assessment of body composition in

- 790 adolescent subjects with anorexia nervosa by bioimpedance. *Medical Engineering & Physics*,
 791 30(6), 783–791.
- 792 Mörk, S., Lackner, S., Müller, W., Gorkiewicz, G., Kashofer, K., Oberascher, A., ... Holasek, S.
 793 (2017). Gut microbiota and body composition in anorexia nervosa inpatients in comparison to
 794 athletes, overweight, obese, and normal weight controls. *International Journal of Eating*
 795 *Disorders*, 50(12), 1421–1431.
- 796 Murray, S. B., Loeb, K. L., & Le Grange, D. (2018). Treatment outcome reporting in anorexia
 797 nervosa: Time for a paradigm shift? *Journal of Eating Disorders*, 6, 10.
- 798 Nakahara, T., Kojima, S., Tanaka, M., Yasuhara, D., Harada, T., Sagiya, K., ... Inui, A. (2007).
 799 Incomplete restoration of the secretion of ghrelin and PYY compared to insulin after food
 800 ingestion following weight gain in anorexia nervosa. *Journal of Psychiatric Research*, 41(10),
 801 814–820.
- 802 Nakai, Y., Hamagaki, S., Takagi, R., Taniguchi, A., & Kurimoto, F. (1999). Plasma concentrations of
 803 tumor necrosis factor- α (TNF- α) and soluble TNF receptors in patients with anorexia
 804 nervosa. *Journal of Clinical Endocrinology and Metabolism*, 84(4), 1226–1228.
- 805 Nieminen, P., Rucker, G., Miettinen, J., Carpenter, J., & Schumacher, M. (2007). Statistically
 806 significant papers in psychiatry were cited more often than others. *Journal of Clinical*
 807 *Epidemiology*, 60(9), 939–946.
- 808 Perry, R. J., & Shulman, G. I. (2018). The role of leptin in maintaining plasma glucose during
 809 starvation. *Postdoc Journal*, 6(3), 3–19.
- 810 Priolella, A., Muscogiuri, G., Sorice, G. P., Lassandro, A. P., Mezza, T., Policola, C., ... Giaccari, A.
 811 (2011). In anorexia nervosa, even a small increase in abdominal fat is responsible for the
 812 appearance of insulin resistance. *Clinical Endocrinology*, 75(2), 202–206.
- 813 Psychogios, N., Hau, D. D., Peng, J., Guo, A. C., Mandal, R., Bouatra, S., ... Wishart, D. S. (2011).
 814 The human serum metabolome. *PloS One*, 6(2), e16957.
- 815 Rigaud, D., Boulieu, A., Tallonneau, I., Brindisi, M. C., & Rozen, R. (2010). Body fluid retention and
 816 body weight change in anorexia nervosa patients during refeeding. *Clinical Nutrition*, 29(6),
 817 749–755.

- 818 Scalfi, L., Marra, M., Caldara, A., Silvestri, E., & Contaldo, F. (1999). Changes in bioimpedance
819 analysis after stable refeeding of undernourished anorexic patients. *International Journal of*
820 *Obesity*, 23(2), 133–137.
- 821 Scalfi, L., Polito, A., Bianchi, L., Marra, M., Caldara, A., Nicolai, E., & Contaldo, F. (2002). Body
822 composition changes in patients with anorexia nervosa after complete weight recovery.
823 *European Journal of Clinical Nutrition*, 56(1), 15–20.
- 824 Schneider, P., Biko, J., Schlamp, D., Trott, G. E., Badura, F., Warnke, A., & Reiners, C. (1998).
825 Comparison of total and regional body composition in adolescent patients with anorexia nervosa
826 and pair-matched controls. *Eating and Weight Disorders*, 3(4), 179–187.
- 827 Schwenk, J. M., Omenn, G. S., Sun, Z., Campbell, D. S., Baker, M. S., Overall, C. M., ... Deutsch, E.
828 W. (2017). The human plasma proteome draft of 2017: Building on the human plasma
829 peptideatlas from mass spectrometry and complementary assays. *Journal of Proteome Research*,
830 16(12), 4299–4310.
- 831 Singhal, V., Tulsiani, S., Campoverde, K. J., Mitchell, D. M., Slattery, M., Schorr, M., ... Klibanski,
832 A. (2018). Impaired bone strength estimates at the distal tibia and its determinants in adolescents
833 with anorexia nervosa. *Bone*, 106, 61–68.
- 834 Solmi, M., Veronese, N., Correll, C. U., Favaro, A., Santonastaso, P., Caregaro, L., ... Stubbs, B.
835 (2016). Bone mineral density, osteoporosis, and fractures among people with eating disorders: A
836 systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*, 133(5), 341–351.
- 837 Suchacki, K. J., & Cawthorn, W. P. (2018). Molecular interaction of bone marrow adipose tissue with
838 energy metabolism. *Current Molecular Biology Reports*, 4(2), 41–49.
- 839 Tagami, T., Satoh, N., Usui, T., Yamada, K., Shimatsu, A., & Kuzuya, H. (2004). Adiponectin in
840 anorexia nervosa and bulimia nervosa. *Journal of Clinical Endocrinology and Metabolism*,
841 89(4), 1833–1837.
- 842 Tanaka, M., Naruo, T., Nagai, N., Kuroki, N., Shiiya, T., Nakazato, M., ... Nozoe, S.-I. (2003).
843 Habitual binge/purge behavior influences circulating ghrelin levels in eating disorders. *Journal*
844 *of Psychiatric Research*, 37(1), 17–22.
- 845 Thompson, S. G., & Sharp, S. J. (1999). Explaining heterogeneity in meta-analysis: A comparison of

methods. *Statistics in Medicine*, 18(20), 2693–2708.

Wan, X., Wang, W., Liu, J., & Tong, T. (2014). Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Medical Research Methodology*, 14, 135.

Watson, H. J., Yilmaz, Z., Thornton, L. M., Hübel, C., Coleman, J. R. I., Gaspar, H. A., ... Bulik, C. M. (2018). Anorexia nervosa genome-wide association study identifies eight loci and implicates metabo-psychiatric origins.

Weinbrenner, T., Züger, M., Jacoby, G. E., Herpertz, S., Liedtke, R., Sudhop, T., ... Berthold, H. K. (2004). Lipoprotein metabolism in patients with anorexia nervosa: a case-control study investigating the mechanisms leading to hypercholesterolaemia. *British Journal of Nutrition*, 91(6), 959–969.

Wells, G. A., Shea, B., O'connell, D., Peterson, J., Welch, V., Losos, M., & Tugwell, P. (2009). The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Retrieved October 1, 2018, from http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp

World Health Organization. (1992). *ICD-10: International Statistical Classification of Diseases and Related Health Problems: 10th revision*. Geneva: World Health Organization.

Yilmaz, Z., Gottfredson, N. C., Zerwas, S. C., Bulik, C. M., & Micali, N. (in press). Developmental premorbid BMI trajectories of adolescents with eating disorders in a longitudinal population cohort. *Journal of the American Academy of Child and Adolescent Psychiatry*.

867 **Table 2** Minimum requirement of variables that should be assessed, reported, and included in
 868 statistical analyses of case-control studies examining anorexia nervosa or other eating
 869 disorders to facilitate reproducibility, meta-analysis, and -regression [adapted from, 111]

| Sampling | Sample characteristics |
|---|---|
| <p>Cases and controls</p> <ul style="list-style-type: none"> • Underlying population: community, hospital • Consecutive sample or selection • If consecutive, attrition & reasons • Diagnosis & ascertainment • Diagnostic schema • Independent validation <p>Controls</p> <ul style="list-style-type: none"> • Repeated measurement • Exclusion of current and history of diagnosis (i.e., screening) • Matching (e.g., age, sex) • Exclusion criteria | <p>Cases and controls</p> <ul style="list-style-type: none"> • Age • Biological sex or gender • Height • Weight • Body mass index • Ancestry • Socioeconomic status <p>Cases</p> <ul style="list-style-type: none"> • Age of onset • Duration of illness |
| Body composition | Menstrual status |
| <ul style="list-style-type: none"> • Fat mass • Fat-free mass • Bone mineral content & density • Ideally: muscle mass | <p>Cases</p> <ul style="list-style-type: none"> • Dysmenorrhea or amenorrhea • Duration of amenorrhea • Age of menarche |

| | |
|---|--|
| <ul style="list-style-type: none"> • Measurement method: MRI, DXA, or BIA • Physical activity (ideally accelerometer data) | <ul style="list-style-type: none"> • If menstruating, stage or day of cycle <p>Controls</p> <ul style="list-style-type: none"> • Stage or day of the menstrual cycle (e.g., follicular phase) |
| Blood sampling | Substances |
| <ul style="list-style-type: none"> • Blood sample type Whole blood, serum, plasma • Fasting state • Fasting period • The time point of blood sampling • Pre-analytics • Storage • Storage duration | <p>Dose and duration of</p> <ul style="list-style-type: none"> • Contraceptives • Supplements & vitamins • Medication <ul style="list-style-type: none"> ○ Prescription ○ Over the counter • Laxatives • Illicit drugs • Alcohol consumption • Smoking behaviour |
| BIA, bioelectrical impedance analysis, DXA, dual-energy X-ray absorptiometry, MRI, magnetic resonance imaging | |

870

871